

Cumulative Index

This index contains all references to methods published in CIPAC 1B, 1C, D, E, F, G, H, J, K, L, M, N and O. All methods from CIPAC 1, CIPAC 1A and from the CIPAC Proceedings 1980 and 1981 have been revised and re-published in CIPAC E (RE's and analytical methods from CIPAC 1), CIPAC F (MT's), and CIPAC H (analytical methods from CIPAC 1A and the Proceedings) as far as they seemed to be still in use. CIPAC code numbers are given in square brackets.

| | CIPAC | page |
|--|-----------------|------------|
| Accelerated storage tests by heating, general methods (MT 46.1) | <i>F</i> | 149 |
| special methods (MT 46.2) | <i>F</i> | 151 |
| combined method (MT 46.3) | <i>J</i> | 128 |
| LN formulations (MT 46.3) | <i>O</i> | 176 |
| Acephate [338], method | <i>H</i> | 5 |
| Acetamiprid [649], method | <i>L</i> | 4 |
| Acetone, anhydrous (RE 34) | <i>E</i> | 286 |
| Acetone soluble material, hot solution (MT 5.1) | <i>F</i> | 21 |
| solution at room temperature (MT 5.2) | <i>F</i> | 22 |
| Acetone insoluble material (MT 27) | <i>F</i> | 88 |
| Acid wash test (MT 79) | <i>F</i> | 210 |
| Adhesion to treated seeds | <i>N</i> | 145 |
| Acidity or alkalinity of formulations (MT 191) | <i>L</i> | 143 |
| Alachlor [204], method, TC, EC, GR, | <i>D</i> | 4 |
| , CS | <i>E</i> | 4 |
| Aldicarb [215], method | <i>H</i> | 10 |
| Aldrin [27], method | <i>B</i> | 1707 |
| <i>d</i> -Allethrin [742], method | <i>L</i> | 16 |
| <i>d</i> -trans-Allethrin (see bioallethrin) | | |
| alpha-Cypermethrin [454], method, TC, WP, EC, SC, UL | | |
| SC (oil enhanced), WG | <i>K</i> | 4 |
| LN | <i>M</i> | 40 |
| alpha HCH, in lindane | <i>G</i> | 106 |
| Alumina, Brockmann activity grade III (MT 50) | <i>F</i> | 156 |
| Ametryn [133], method, GC | <i>H</i> | 22 |
| Amidate, method | <i>E</i> | 142 |
| Amines, identification of (MT 152) | <i>F</i> | 357 |
| Aminocarb [293] | <i>D</i> | 9 |
| 2-Aminopyridine, preparation of complexes of nitro compounds (MT 134) | <i>F</i> | 328 |

CUMULATIVE INDEX

| | CIPAC | page |
|--|-----------|------------|
| Amisulbrom [789], method, TC, SC, WG | O | 4 |
| Amitraz [362], method | <i>G</i> | 5 |
| Amitrole [90], method, | | |
| potentiometric | <i>IB</i> | 1720 |
| argentometric | <i>IB</i> | 1722 |
| HPLC | <i>J</i> | 4 |
| Amitrole + monuron [90 + 99] | <i>IB</i> | 1725 |
| Amitrole + simazin [90 + 22] | <i>IB</i> | 1725 |
| Amitrole + NH ₄ CNS | <i>J</i> | 9 |
| Amitrole + diuron [90 + 100] | <i>J</i> | 8 |
| Amitrole + terbuthylazine [90 + 234] | <i>J</i> | 9 |
| Ammonia, solutions (RE 1) | <i>E</i> | 245 |
| Ammonia-ammonium chloride buffer solution-pH 10 (MT 107) | <i>F</i> | 261 |
| Ammonium cerium (IV) nitrate (RE 148) | <i>E</i> | 383 |
| Ammonium hexanitratocerate (IV) (RE 148) | <i>E</i> | 383 |
| diAmmonium hydrogen citrate (RE 91) | <i>E</i> | 362 |
| Ammonium iron (III) sulfate, indicator solution (RE 2) | <i>E</i> | 246 |
| diAmmonium iron (II) sulfate, standard solutions (RE 38) | <i>E</i> | 297 |
| diAmmonium peroxosulfate (RE 132) | <i>E</i> | 375 |
| Ammonium thiocyanate, standard solution (RE 3) | <i>E</i> | 249 |
| Ammonium trioxovanadate (V), solution (RE 31) | <i>E</i> | 283 |
| Anilazine [294], method | <i>E</i> | 5 |
| Aromatic solvent naphtha (RE 62) | <i>E</i> | 328 |
| Arsenic, determination of (MT 99) | <i>F</i> | 251 |
| Arsenic in sulfur, method | <i>E</i> | 205 |
| Arsenic in zineb, method | <i>E</i> | 231 |
| Arsenic in ziram, method | <i>E</i> | 237 |
| Arsenic (III) oxide, standard solution (RE 4) | <i>E</i> | 250 |
| Asulam [240], method | <i>H</i> | 26 |
| Atrazine [91], method, GC | <i>H</i> | 37 |
| Attrition of tablets (MT 193) | O | 204 |
| replaces MT 193 Handbook L, p. 147 | | |
| Attrition resistance of granules (MT 178) | <i>H</i> | 302 |
| Attrition resistance of dispersible granules (MT 178.2) | <i>K</i> | 140 |
| Azadirachtin [627], method | <i>M</i> | 4 |
| Azimsulfuron [584], method | <i>L</i> | 23 |
| Azinphos-methyl [37], method, HPLC | <i>E</i> | 12 |
| IR | <i>IC</i> | 1970 |
| Azoxystrobin [571], method | <i>M</i> | 10 |
| Barium hydroxide, standard solutions (RE 5) | <i>E</i> | 251 |
| Bendiocarb [232], method | <i>D</i> | 10 |

CUMULATIVE INDEX

| | CIPAC | page |
|--|-----------------|-----------|
| Benfluralin [285], method, GC | <i>IB</i> | 1728 |
| UV | <i>IB</i> | 1726 |
| Benfuracarb [501], method | <i>H</i> | 38 |
| Benomyl [206], method | <i>D</i> | 14 |
| Bensulfuron-methyl [502], method | <i>K</i> | 6 |
| Bensultap [464], method | <i>E</i> | 17 |
| Bentazone [366], method | <i>IC</i> | 1973 |
| Benzoylprop-ethyl [229], method | <i>IB</i> | 1732 |
| Benzoylprop-ethyl, pure (PP 229.a) | <i>IB</i> | 1930 |
| beta-Cyfluthrin [482], method | <i>H</i> | 43 |
| BHC (see HCH) | | |
| BHTO in fenbutatin oxide | <i>D</i> | 81 |
| Bifenox [413], method | <i>G</i> | 11 |
| Binapacryl [138], method | <i>D</i> | 67 |
| Bioallethrin [203], method, TC | <i>H</i> | 52 |
| TC, LV | <i>L</i> | 31 |
| Bioallethrin + piperonyl butoxide [203 + 33] | <i>H</i> | 55 |
| S-Bioallethrin [750], method TC, LV | <i>L</i> | 38 |
| EW | <i>M</i> | 18 |
| 2,2'-Bipyridyl (RE 61) | <i>E</i> | 326 |
| 2,2'-Bipyridyl in diquat | <i>H</i> | 161 |
| 4,4'-Bipyridyl (RE 60) | <i>E</i> | 325 |
| 4,4'-Bipyridyl in paraquat | <i>H</i> | 216 |
| Bis (ethylenediamine) copper (II) sulfate (RE 81) | <i>E</i> | 359 |
| Bitertanol [386], method | <i>J</i> | 14 |
| Borax (RE 33) | <i>E</i> | 376 |
| Boscalid [673], method | <i>N</i> | 4 |
| Brodifacoum [370], chiral HPLC method, TC, RB | <i>O</i> | 12 |
| HPLC method | <i>IC</i> | 1981 |
| Bromadiolone [371], method | <i>E</i> | 23 |
| Bromacil [139], method | <i>IC</i> | 1986 |
| Bromocresol green, solutions (RE 55) | <i>E</i> | 317 |
| Bromophenol blue, solution (RE 6) | <i>E</i> | 254 |
| Bromophos [5], method | <i>IB</i> | 1738 |
| Bromothymol blue, solutions (RE 7) | <i>E</i> | 254 |
| Bromoxynil [87], method | <i>IC</i> | 1989 |
| pure (PP 87) | <i>IC</i> | 2326 |
| - octanoate, GC method 1, TC | <i>IC</i> | 1998 |
| GC method 2, TC,EC | <i>IC</i> | 2004 |
| pure (PP 87.oct) | <i>IC</i> | 2330 |
| - heptanoate | <i>H</i> | 57 |

CUMULATIVE INDEX

| | CIPAC | page |
|---|-----------------|-----------|
| Buffer Solutions | | |
| Phosphate buffer (MT 19) | <i>F</i> | 69 |
| Bulk density (MT 186) | <i>K</i> | 151 |
| Bupirimate, identity test (MT 164) | <i>F</i> | 406 |
| Bupirimate [261], method | <i>IC</i> | 2007 |
| Bupirimate, pure (PP 261) | <i>IC</i> | 2340 |
| Butachlor [354], method | <i>D</i> | 17 |
| Butylate [266], method | <i>IB</i> | 1744 |
| Butylmagnesium chloride (RE 139) | <i>E</i> | 379 |
| | | |
| Cadmium, standard solution (RE 131) | <i>E</i> | 374 |
| Captan [40], method, GC, | <i>IC</i> | 2015 |
| HPLC | <i>IC</i> | 2013 |
| Carbaryl [26], method | <i>M</i> | 19 |
| Carbendazim [263], method, UV | <i>H</i> | 61 |
| HPLC | <i>H</i> | 62 |
| Carbetamide [95], method | <i>E</i> | 28 |
| Carbon disulfide (RE 153) | <i>E</i> | 384 |
| Carbofuran [276], method | <i>D</i> | 20 |
| Carbosulfan [417], method | <i>E</i> | 35 |
| Cartap [387], method | <i>D</i> | 24 |
| Cerium (IV) sulfate, standard solutions (RE 99) | <i>E</i> | 368 |
| Cetylpyridinium bromide, solution (RE 11) | <i>E</i> | 258 |
| Chinomethionat [172], method | <i>IC</i> | 2019 |
| Chloramben[141], method | <i>IB</i> | 1747 |
| Chlordimeform [277], method | <i>D</i> | 28 |
| Chlorfenapyr [570], method, TC, SC | <i>O</i> | 22 |
| Chlorfenvinphos [88], method | <i>H</i> | 67 |
| Chloridazon [111], method | <i>D</i> | 31 |
| water dispersible granules | <i>H</i> | 76 |
| Chlorides, in phenoxyalkanoic acids (MT 68.1) | <i>F</i> | 195 |
| Chlorides, test for (MT 117) | <i>F</i> | 282 |
| Chlorides, total chlorides (MT 100) | <i>F</i> | 254 |
| Chlormequat [143], titration method | <i>D</i> | 39 |
| ion chromatographic method | <i>H</i> | 77 |
| Chlorobenzilate [119], method | <i>H</i> | 81 |
| 4-Chloro- <i>m</i> -cresol (RE 104) | <i>E</i> | 371 |
| 4-Chloro- <i>o</i> -cresol (RE 109) | <i>E</i> | 372 |
| 6-Chloro- <i>p</i> -cresol (RE 68) | <i>E</i> | 335 |
| 1-Chloro-2,3-epoxyethane (MT 77) | <i>F</i> | 207 |
| Chloroform (RE 64) | <i>E</i> | 330 |
| Chloroform solubles, hot solutions (MT 87.1) | <i>F</i> | 221 |

CUMULATIVE INDEX

| | CIPAC | page |
|--|-----------------|------------------|
| cold solutions (MT 87.2) | <i>F</i> | 222 |
| 6-Chloro-2-methylphenoxyacetic acid (RE 65) | <i>E</i> | 332 |
| γ-Chlorophosphamidon, method | <i>E</i> | 182 |
| Chloropropylate [120], method | <i>H</i> | 85 |
| Chlorothalonil [288], method | <i>K</i> | 13 |
| Chlorothalonil + HCB [288 +161], method | <i>K</i> | 13 |
| 6-Chloro- <i>o</i> -tolylloxyacetic acid (RE 65) | <i>E</i> | 332 |
| Chloroxuron [187], method | <i>1B</i> | 1751 |
| Chlorphoxim, [341], method | <i>D</i> | 24 |
| Chlorpropham [43], method | <i>1B</i> | 1757 |
| Chlorpropham + propham [43 + 63] | <i>1C</i> | 2025 |
| Chlorpyrifos [221], method, TC, DP, WP, EC, GR | <i>1C</i> | 2028 |
| UL | <i>K</i> | 23 |
| Chlorsulfuron [441], method, TC, WG | <i>H</i> | 89 |
| WP | <i>K</i> | 24 |
| Chlorthal [328], method, GC | <i>1C</i> | 2032 |
| IR | <i>1C</i> | 2034 |
| CIPAC method | <i>O</i> | <i>1</i> |
| Clodinafop [683]; method | <i>M</i> | 26 |
| Clofentezine [418], method | <i>G</i> | 18 |
| Clothianidin [738], method | <i>N</i> | 14 |
| Conductivity, determination of (MT 32) | <i>F</i> | 103 |
| Congo red, solution (RE 9) | <i>E</i> | 256 |
| Copper [44], method | <i>E</i> | 42 |
| Copper ethylenediamine sulfate (RE 81) | <i>E</i> | 350 |
| Copper, water soluble | | |
| colorimetric method with bathocuproine (98.1) | <i>F</i> | 246 |
| atomic absorption method (MT 98.2) | <i>F</i> | 248 |
| Copper + mancozeb [44 + 34], method | <i>H</i> | 96 |
| Copper + maneb [44 + 61], method | <i>H</i> | 100 |
| Copper + zineb [44 + 25], method | <i>H</i> | 100 |
| Copper + ziram [44 + 31], method | <i>H</i> | 101 |
| Correction of interfering peaks (MT 114) | <i>F</i> | 278 |
| Coumatetralyl [189], method | <i>G</i> | 24 |
| <i>m</i> -Cresol (RE 100) | <i>E</i> | 370 |
| <i>o</i> -Cresol (RE 70) | <i>E</i> | 338 |
| <i>p</i> -Cresol (RE 101) | <i>E</i> | 370 |
| Crystal violet, solution (RE 129) | <i>E</i> | 373 |
| Cyanazine [230], method | <i>1C</i> | 2039 |
| Cyanazine, pure (PP 230) | <i>1C</i> | 2335 |
| Cyazofamid, [653], method, TC, SC | <i>O</i> | <i>31</i> |

CUMULATIVE INDEX

| | CIPAC | page |
|--|-----------|------|
| Cycloate [214], method | <i>H</i> | 102 |
| Cyclohexane (RE 155) | <i>E</i> | 386 |
| Cycloxydim [510], method | <i>K</i> | 25 |
| Cyfluthrin [385], method | <i>H</i> | 106 |
| beta-Cyfluthrin [482], method | <i>H</i> | 43 |
| lambda-Cyhalothrin [463], method, TC, WP, EC, UL | <i>E</i> | 49 |
| LN | <i>N</i> | 83 |
| lambda-Cyhalothrin, free, determination in CS formulations (MT 189) | <i>L</i> | 137 |
| lambda-Cyhalothrin, determination of release properties CS formulations (MT 190) | <i>L</i> | 140 |
| Cyhexatin [289], method | <i>D</i> | 46 |
| Cymoxanil [419], method | <i>J</i> | 22 |
| Cypermethrin, identity test (MT 163) | <i>F</i> | 404 |
| Cypermethrin [332], method, GC | <i>1C</i> | 2047 |
| diastereomeric ratio, HPLC | <i>1C</i> | 2052 |
| alpha-Cypermethrin [454], method, TC, WP, EC, SC, UL | <i>H</i> | 14 |
| SC (oil enhanced), WG | <i>K</i> | 4 |
| LN | <i>M</i> | 40 |
| Cyphenothrin [804], method | <i>M</i> | 44 |
| Cyproconazole [600], method | <i>J</i> | 29 |
| Cyprodinil [511], method | <i>N</i> | 25 |
| Cyromazin [420], method | <i>M</i> | 54 |
| 2,4-D [1], method, titration method | <i>1C</i> | 2060 |
| GC | <i>1C</i> | 2061 |
| HPLC | <i>1C</i> | 2062 |
| -salts titration method | <i>1C</i> | 2063 |
| GC | <i>1C</i> | 2065 |
| HPLC | <i>1C</i> | 2065 |
| -esters titration method | <i>1C</i> | 2068 |
| GC | <i>1C</i> | 2069 |
| HPLC | <i>1C</i> | 2071 |
| 2,4-D + dicamba [1+85] (salts), GC | <i>1C</i> | 2071 |
| IR | <i>H</i> | 133 |
| 2,4-D + dicamba + mecoprop [1 + 85 + 51] (salts) | <i>D</i> | 51 |
| 2,4-D + picloram [1 + 174] | <i>1B</i> | 1757 |
| 2,4-D + 2,4,5-T [1 + 6](esters) | <i>1C</i> | 2072 |
| 2,4-D, pure (PP 1) | <i>1C</i> | 2307 |
| Dalapon [52], method (Na + Mg salt) | <i>D</i> | 54 |
| Dazomet [146], method, CS ₂ method | <i>1C</i> | 2074 |
| HPLC | <i>J</i> | 37 |

CUMULATIVE INDEX

| | CIPAC | page |
|--|-----------------|------------|
| 2,4-DB [83], method | <i>IC</i> | 2078 |
| -K salt, titration method | <i>IC</i> | 2080 |
| GC | <i>IC</i> | 2082 |
| 2,4-DB, pure (PP 83) | <i>IC</i> | 2321 |
| DDT [3], method | <i>E</i> | 58 |
| DDT + endrin [3 + 28] | <i>IB</i> | 1760 |
| Degree of dissolution and solution stability (MT 179.1) | <i>O</i> | 189 |
| replaces MT 179 | | |
| Deltamethrin [333], method, TC, WP, EC, UL, DP | <i>D</i> | 57 |
| method, TC, WG, WP, EC, SC, DP, EG, EW, UL | <i>L</i> | 45 |
| TB, LN | <i>M</i> | 65 |
| PE LN | <i>N</i> | 34 |
| PE + PP LN incorporated | <i>O</i> | 43 |
| method, isomers, TC | <i>O</i> | 39 |
| Desmetryn [147], method, GC | <i>IB</i> | 1766 |
| titration method | <i>IB</i> | 1765 |
| Deniges reagent (RE 57) | <i>E</i> | 319 |
| Deschlorophosphamidon, method | <i>E</i> | 182 |
| Detection and identification of impurities in substituted phenylurea herbicides (MT 142) | <i>F</i> | 336 |
| Determination of organochlorine pesticides in aqueous emulsions (MT 65) | <i>F</i> | 190 |
| Determination of pH values, general method (MT 75.1) | <i>F</i> | 205 |
| pH of aqueous dispersions (MT 75.2) | <i>F</i> | 206 |
| pH of diluted and undiluted aqueous solutions (MT 75.3) | <i>J</i> | 131 |
| Determination of seed-to-seed uniformity of distribution for liquid seed-treatment formulations (MT 175) | <i>F</i> | 438 |
| Diazinon [15], method | <i>H</i> | 122 |
| Diazomethane (RE 35) | <i>E</i> | 286 |
| 1,2-Dibromoethane [125], method | <i>IC</i> | 2115 |
| Dicamba [85], method, infrared method | <i>H</i> | 127 |
| HPLC method | <i>K</i> | 32 |
| Dicamba + 2,4-D [85 + 1], method | <i>H</i> | 133 |
| Dicamba + MCPA [85 + 2], method | <i>H</i> | 131 |
| Dicamba + 2,4-D + mecoprop [85 + 1 + 51] (salts) | <i>D</i> | 51 |
| Dichlobenil [73], method, GC(cap) | <i>E</i> | 65 |
| GC | <i>IB</i> | 1769 |
| Dichlofluanid [74], method, titration method | <i>IB</i> | 1774 |
| HPLC | <i>IC</i> | 2083 |
| 2,4-Dichloroanisole in bifenoX | <i>G</i> | 14 |
| 2,4-Dichlorobenzoic acid (RE 79) | <i>E</i> | 347 |

CUMULATIVE INDEX

| | CIPAC | page |
|--|-------------------------|-------------|
| 4,6-Dichloro- <i>o</i> -cresol (RE 69) | <i>E</i> | 335 |
| Dichlorodifluoromethane insoluble material (MT 16) | <i>F</i> | 52 |
| 4,6-Dichloro-2-methylphenoxyacetic acid (RE 66) | <i>E</i> | 333 |
| 2,4-Dichlorophenol in bifenox | <i>G</i> | 14 |
| 2,6-Dichlorophenol (RE 72) | <i>E</i> | 339 |
| 4,6-Dichloro- <i>o</i> -tolylloxyacetic acid (RE 66) | <i>E</i> | 333 |
| Dichlorprop [84], method, titration method | <i>IC</i> | 2087 |
| | GC | 2087 |
| | HPLC | 2088 |
| | HPLC | 32 |
| -salts titration method | <i>IC</i> | 2092 |
| | GC | 2093 |
| | HPLC | 2093 |
| Dichlorprop, pure (PP 84) | <i>IC</i> | 2323 |
| Dichlorprop-P [476], method | <i>G</i> | 32 |
| Dichlorvos [11], method,GC | <i>H</i> | 135 |
| Diclofop [358], method | <i>IC</i> | 2096 |
| Dicofol [123], method | <i>D</i> | 67 |
| Dieldrin [16], method | <i>IB</i> | 1781 |
| Diethyl ether (RE 12) | <i>E</i> | 258 |
| Diflovidazin [734], method | <i>L</i> | 61 |
| Di flubenzuron [339], method, TC,WP, | <i>H</i> | 141 |
| | TC,WP (revised methods) | <i>N</i> 38 |
| | SC,GR,TB | <i>N</i> 42 |
| Di flufenican [462], method | <i>H</i> | 147 |
| Dihydrosafrole in piperonyl butoxide, TC | <i>O</i> | 104 |
| Diisopropyl ether (RE 63) | <i>E</i> | 328 |
| Dilute emulsion stability (MT 20) | <i>F</i> | 71 |
| colorimetric method (MT 173) | <i>F</i> | 431 |
| Dilution stability (MT 41) | <i>F</i> | 131 |
| Dilution stability of aqueous solutions (MT 41.1) | <i>O</i> | 174 |
| Disintegration of tablets (MT 197) | <i>O</i> | 212 |
| Dissolution degree and solution stability (MT 179) | <i>H</i> | 307 |
| Dissolution degree and solution stability (MT 179.1) | <i>O</i> | 189 |
| replaces MT 179 | | |
| Dimedone derivative (MT 28) | <i>F</i> | 89 |
| Dimercury dicyanide oxide (RE 154) | <i>E</i> | 383 |
| Dimethoate [59], HPLC method | <i>E</i> | 69 |
| Dimethoate [59], GLC method | <i>H</i> | 153 |
| Dimethomorph [483], method | <i>G</i> | 32 |
| <i>O,O</i> -Dimethyl S-methylcarbamoylmethylphosphorodithioate (RE 86) | <i>E</i> | 359 |

CUMULATIVE INDEX

| | CIPAC | page |
|---|-----------------|------------|
| <i>O,O</i> -Dimethyl methylphosphoramidothioate, method | <i>E</i> | 142 |
| <i>O,S</i> -Dimethyl methylphosphoramidothioate, method | <i>E</i> | 142 |
| <i>O,S</i> -Dimethyl 4-nitrophenylphosphorothioate, method | <i>E</i> | 169 |
| <i>O,O</i> -Dimethyl phosphoramidothioate, method | <i>E</i> | 142 |
| <i>O,O</i>-dimethyl phosphorochloridothioate | <i>O</i> | 118 |
| Dimethyl sulfate (RE 13) | <i>E</i> | 259 |
| Dimoxystrobin [739], method, TC, SE, SC | <i>O</i> | 47 |
| Dinotefuran [749], method | <i>L</i> | 67 |
| Dinoterb [403], method | <i>IB</i> | 1797 |
| Dinitro compounds, solubility of salts and alkali insolubles (MT 108) | <i>F</i> | 262 |
| mineral acidity of (MT 109) | <i>F</i> | 263 |
| Dioxane (RE 49) | <i>E</i> | 311 |
| Diquat [55], method | <i>E</i> | 74 |
| identity tests | <i>G</i> | 47 |
| free 2,2'-bipyridyl | <i>H</i> | 161 |
| Terpyridines in | <i>M</i> | 68 |
| Diquat dibromide (PP 55) | <i>E</i> | 387 |
| Dispersion stability of suspo-emulsions (MT 180) | <i>H</i> | 310 |
| SE, CD and OD formulations | <i>O</i> | 192 |
| Dissolution rate of water-soluble bags (MT 176) | <i>F</i> | 440 |
| Disulfoton [152], method | <i>IC</i> | 2101 |
| Dithianon [153], method, HPLC | <i>G</i> | 50 |
| colorimetric | <i>IC</i> | 2105 |
| Dithiocarbamates, determination of anions (MT 154) | <i>F</i> | 361 |
| Identity test for certain alkylenebis(dithiocarbamates) | <i>F</i> | 320 |
| Qualitative procedure for confirmation of the presence of a dithiocarbamate or thiuram disulfide (MT 153) | <i>F</i> | 360 |
| Dithizone (RE 59) | <i>E</i> | 371 |
| Dodine [101], method | <i>IB</i> | 1802 |
| Dustability test after accelerated storage (MT 34) | <i>F</i> | 107 |
| Dust content and apparent density of granular formulations, | | |
| sampling (MT 58.1) | <i>F</i> | 173 |
| preparation of the sample (MT 58.2) | <i>F</i> | 173 |
| apparent density after compaction without pressure (MT 58.3) | <i>F</i> | 175 |
| Dustiness of granular products (MT 171) | <i>F</i> | 425 |
| Edifenphos [409], method | <i>E</i> | 79 |
| EDTA sodium salt, solutions (RE 37) | <i>E</i> | 295 |
| Emulsion characteristics of emulsifiable concentrates, | | |
| five per cent v/v oil phase when diluted (MT 36.1) | <i>F</i> | 108 |
| dispersion stability of aqueous emulsions (MT 36.2) | <i>F</i> | 112 |
| emulsion characteristics and re-emulsification properties | | |

CUMULATIVE INDEX

| | CIPAC | page |
|---|-----------------|------------|
| Fenbutatin oxide [359], method | <i>D</i> | 77 |
| Fenhexamid [603], method | <i>M</i> | 80 |
| Fenitrothion [35], method, TC, method from CIPAC 1C p 2117 was replaced | <i>N</i> | 46 |
| Fenobucarb [390], method, TLC | <i>D</i> | 87 |
| HPLC | <i>D</i> | 89 |
| Fenoprop [118], method, titration method | <i>1C</i> | 2121 |
| GC | <i>1C</i> | 2122 |
| salts, titration method | <i>1C</i> | 2122 |
| GC | <i>1C</i> | 2123 |
| Fenoprop, pure (PP 118) | <i>1C</i> | 2333 |
| Fenoxaprop-P [484], method, TC, EW | <i>J</i> | 51 |
| TC, EW, EC, SE, OD | <i>M</i> | 89 |
| Fenpropimorph [427], method | <i>G</i> | 62 |
| Fensulfothion [303], method, HPLC | <i>1C</i> | 2126 |
| GC | <i>D</i> | 92 |
| Fenthion [79], method, col. | <i>1B</i> | 1830 |
| method, GC, TC, WP, GR, EC, EW | <i>L</i> | 81 |
| UL, DP | <i>M</i> | 96 |
| Fentin acetate [489], method | <i>E</i> | 89 |
| Fentin acetate + maneb [489 + 61], GC | <i>1B</i> | 1837 |
| HPLC | <i>E</i> | 89 |
| Fentin hydroxide [490], method | <i>E</i> | 94 |
| Fentin hydroxide + maneb [490 + 61] GC | <i>1B</i> | 1840 |
| HPLC | <i>E</i> | 93 |
| Fenvalerate, identity test (MT 163) | <i>F</i> | 404 |
| Fenvalerate [334], method, TC | <i>D</i> | 101 |
| EC | <i>G</i> | 68 |
| Ferroun sulfate, solution (RE 149) | <i>E</i> | 383 |
| Fipronil [581], method TC, EC, UL | <i>J</i> | 60 |
| FS, SC, WG, GR, FG | <i>N</i> | 60 |
| Flash point, Abel method (MT 12.1) | <i>F</i> | 31 |
| Tag closed tester (MT 12.2) | <i>F</i> | 37 |
| Pensky-Martens closed tester (MT 12.3) | <i>F</i> | 40 |
| Flazasulfuron [595], method, TC, WG | <i>O</i> | 60 |
| Flowability, of powders (MT 44) | <i>F</i> | 145 |
| of water dispersible granules (MT 172) | <i>F</i> | 430 |
| of granular preparations after accelerated storage under pressure (MT 172.1) | <i>O</i> | 187 |
| Fluazifop-butyl [395], method | <i>D</i> | 106 |
| Fluazifop-P-butyl [467], method | <i>G</i> | 71 |
| Fluazinam [521], method, TC, SC | <i>O</i> | 68 |

CUMULATIVE INDEX

| | CIPAC | page |
|---|-----------------|------------|
| Flufenoxuron [470], method | <i>M</i> | 98 |
| Flumioxazin [578], method, TC, WP | <i>O</i> | 77 |
| Fluometuron [159], method | <i>IB</i> | 1842 |
| Flusilazole [435], method | <i>H</i> | 171 |
| Folpet [75], method | <i>IB</i> | 1845 |
| Foam persistent, general method (MT 47.3) | <i>O</i> | 177 |
| Foaming, persistent (MT 47.1) | <i>F</i> | 152 |
| of suspension concentrates (MT 47.2) | <i>F</i> | 152 |
| Formaldehyde [156], method | <i>IC</i> | 2130 |
| Formothion [160], method | <i>H</i> | 178 |
| Fosetyl [384], method, titration method | <i>G</i> | 82 |
| HPLC | <i>J</i> | 66 |
| Fosthiazate [585], method, TC, GR | <i>O</i> | 83 |
| Free acidity or alkalinity, methyl red indicator method (MT 31.1) | <i>F</i> | 96 |
| electrometric procedure (MT 31.2) | <i>F</i> | 98 |
| acidity of petroleum products (MT 31.3) | <i>F</i> | 99 |
| Free acidity or alkalinity of formulations (MT 191) | <i>L</i> | 143 |
| Free acidity of phenoxyalkanoic esters (MT 66) | <i>F</i> | 193 |
| Free amines in urea herbicides (MT 141) | <i>F</i> | 333 |
| Free 2,2'-bipyridyl in diquat | <i>H</i> | 161 |
| Free 4,4'-bipyridyl in paraquat | <i>H</i> | 216 |
| Free lambda-cyhalothrin in CS formulations (MT 189) | <i>L</i> | 137 |
| Free pirimiphos-methyl in CS formulations (MT 189.2) | <i>O</i> | 193 |
| Free parathion-methyl in CS formulations (MT 188) | <i>L</i> | 132 |
| Free phenols in phenoxyalkanoic herbicides, | | |
| GLC method (MT 69) | <i>F</i> | 197 |
| HPLC method (MT 155) | <i>F</i> | 362 |
| Freezing mixtures, (5 ± 1°C) (MT 14.1) | <i>F</i> | 44 |
| (10 ± 1°C) (MT 14.2) | <i>F</i> | 44 |
| Freezing point (MT 1) | <i>F</i> | 3 |
| Friability of tablets (MT 193) | <i>L</i> | 147 |
| replaced by Attrition of tablets MT 193 | <i>O</i> | 204 |
| Gas chromatography of phenoxyalkanoic acids and other herbicides (MT 129) | <i>F</i> | 292 |
| Glufosinate [437], method | <i>G</i> | 89 |
| Glyphosate [284], method, TC, SL | <i>IC</i> | 2132 |
| SG | <i>H</i> | 182 |
| Haloxyfop-P-methyl [526], method | <i>N</i> | 67 |
| Hardness of water (MT 73) | <i>F</i> | 201 |
| Total hardness of water (MT 73.1) | <i>O</i> | 179 |

CUMULATIVE INDEX

| | CIPAC | page |
|---|-----------|------|
| HCB (in chlorothalonil) [161, (288)] | <i>K</i> | 13 |
| HCH (lindane in) [4, (488)], method | <i>IC</i> | 1977 |
| HEOD (see dieldrin) | | |
| Heptane (RE 50) | <i>E</i> | 313 |
| Heptane-insoluble material (MT 101) | <i>F</i> | 255 |
| Hexadecanoic acid (RE 80) | <i>E</i> | 348 |
| Hexane (RE 76) | <i>E</i> | 344 |
| Hexazinone [374], method | <i>J</i> | 71 |
| HHDN (see aldrin) | | |
| Hydrochloric acid, standard solutions (RE 14) | <i>E</i> | 262 |
| Hydrogen peroxide (RE 56) | <i>E</i> | 318 |
| Hydrogen sulfide and thiols (MT 78) | <i>F</i> | 208 |
| Hydrolysable organic chlorine, | | |
| DDT residues of technical material (MT 64.4) | <i>F</i> | 187 |
| DDT dusts and dispersible powders (MT 64.5) | <i>F</i> | 189 |
| DDT emulsifiable concentrates and solutions (MT 64.6) | <i>F</i> | 189 |
| HCH and lindane, dried residues, or technical materials (MT 64.1) | <i>F</i> | 183 |
| HCH or lindane dusts and dispersible powders (MT 64.2) | <i>F</i> | 186 |
| HCH or lindane emulsifiable concentrates and solutions (MT 64.3) | <i>F</i> | 187 |
| Hydroxylammonium chloride (RE 74) | <i>E</i> | 350 |
| Icaridin [740], method | <i>K</i> | 64 |
| Identification of amines (MT 152) | <i>F</i> | 357 |
| Identification of urea herbicides (MT 137) | <i>F</i> | 329 |
| Identity test for certain dithiocarbamates and thiram and their mixtures (MT 130) | <i>F</i> | 321 |
| Identity tests for permethrin, cypermethrin and fenvalerate (MT 163) | <i>F</i> | 404 |
| Identity tests for pirimicarb, bupirimate, ethirimol, pirimiphos-ethyl and pirimiphos-methyl (MT 164) | <i>F</i> | 406 |
| Ignition tests, assessment of spontaneous ignition potential of dithio-carbamates (MT 84.1) | <i>F</i> | 218 |
| Imazalil [335], method | <i>E</i> | 95 |
| Imidacloprid [582], method, TC, WP, SC, GR, fertilizer formulations | <i>H</i> | 185 |
| TC, WG, SC, WS, FS | <i>K</i> | 70 |
| SL, OD | <i>L</i> | 89 |
| Indoxacarb, [612], method | <i>N</i> | 73 |
| Iodides, test for (MT 118) | <i>F</i> | 283 |
| Iodine, standard solutions (RE 16) | <i>E</i> | 264 |
| Iodine, monochloride, solution (RE 58) | <i>E</i> | 320 |

CUMULATIVE INDEX

| | CIPAC | page |
|--|-----------------|------------|
| Ioxynil [86], method | <i>E</i> | 101 |
| Ioxynil octanoate [86.407], method | <i>G</i> | 94 |
| Iprobenfos [393], method | <i>D</i> | 110 |
| Iprodione [278], method, TC, WP, SC | <i>G</i> | 98 |
| WG | <i>L</i> | 90 |
| IR 3535 see Ethyl butylacetamidopropionate | | |
| Iron, determination of (MT 95) | <i>F</i> | 236 |
| Iron (III) chloride (RE 84) | <i>E</i> | 355 |
| Isobutyl benzoate (RE 92) | <i>E</i> | 362 |
| Isochloridazone | <i>D</i> | 35 |
| Isofenphos [412], method | <i>D</i> | 114 |
| Isolation of active ingredient, extraction with acetone (MT 37.1) | <i>F</i> | 115 |
| extraction with light petroleum (MT 37.2) | <i>F</i> | 115 |
| Isomalathion, method | <i>IB</i> | 1855 |
| Isopirimiphos-methyl in pirimiphos-methyl | <i>O</i> | 116 |
| Isoprocarb [396], method | <i>D</i> | 118 |
| Isoprothiolane [456], method | <i>E</i> | 105 |
| Isoproturon [336], method | <i>E</i> | 110 |
| | | |
| John Innes Potting compost for soil testing of seed germination (MT 26) | <i>F</i> | 87 |
| | | |
| Kerosene (RE 73) | <i>E</i> | 342 |
| Kerosene insoluble material (MT 8) | <i>F</i> | 25 |
| Kieselguhr, for GLC (MT 86.1) | <i>F</i> | 219 |
| for partition chromatography (MT 86.2) | <i>F</i> | 220 |
| Kresoxim-methyl [568], method | <i>K</i> | 77 |
| | | |
| lambda-Cyhalothrin [463], method, TC, WP, EC, UL | <i>E</i> | 49 |
| CS | <i>K</i> | 86 |
| LN | <i>N</i> | 83 |
| Lead, dithizone general method (MT 92.1) | <i>F</i> | 224 |
| dithizone method B (MT 92.2) | <i>F</i> | 229 |
| Lead, standard solution (RE 90) | <i>E</i> | 361 |
| Lindane [488], method, SC | <i>G</i> | 105 |
| TC, WP | <i>1C</i> | 2135 |
| - alpha HCH | <i>G</i> | 108 |
| - in HCH [4] | <i>1C</i> | 1979 |
| Liquid paraffin (RE 8) | <i>E</i> | 255 |
| Loss on drying, oven drying at 60°C (MT 17.1) | <i>F</i> | 55 |
| vacuum drying at 45°C, 24 h (MT 17.2) | <i>F</i> | 56 |
| over drying at 100°C at constant weight (MT 17.3) | <i>F</i> | 57 |

CUMULATIVE INDEX

| | CIPAC | page |
|---|-----------|------|
| Mecoprop + 2,4-D + dicamba [51 + 1 + 8](salts) | <i>D</i> | 51 |
| Mecoprop, pure (PP 51) | <i>1C</i> | 2319 |
| Mecoprop-P [475], method | <i>G</i> | 109 |
| Mefenpyr-diethyl, [651], method | <i>N</i> | 87 |
| Melting point (MT 2) | <i>F</i> | 5 |
| Mepiquat [440], method | <i>J</i> | 78 |
| Mercurial impurities in technical and formulated mercury compounds (MT 110) | <i>F</i> | 264 |
| Mercury, determination on treated seed (MT 158) | <i>F</i> | 387 |
| Mercuric oxycyanide (RE 154) | <i>E</i> | 385 |
| Mercury(II)chloride (RE 135) | <i>E</i> | 378 |
| Mercury(II) salts, characteristic reactions of (MT 116) | <i>F</i> | 280 |
| Mercury(II)sulfate, solution (RE 57) | <i>E</i> | 319 |
| Metalaxyl [365], method | <i>E</i> | 123 |
| Metamitron [381], method | <i>D</i> | 124 |
| Metam-sodium [20], method | <i>E</i> | 131 |
| Metazachlor [411], method | <i>E</i> | 134 |
| Methabenzthiazuron [201], method | <i>G</i> | 116 |
| Methamidophos [355], method | <i>E</i> | 139 |
| Methanol (RE 47) | <i>E</i> | 309 |
| Methiocarb [165], method | <i>D</i> | 130 |
| Methomyl [264], method | <i>H</i> | 199 |
| Methoprene [414], method | <i>M</i> | 115 |
| Methoprotryne [94], method | <i>1B</i> | 1864 |
| <i>N</i> -Methyl amidate, method | <i>E</i> | 142 |
| Methylcyclopropene [767], method | <i>N</i> | 99 |
| Methyl (dimethoxyphosphinothioylthio) acetate (RE 88) | <i>E</i> | 360 |
| Methyl-methamidophos, method | <i>E</i> | 142 |
| Methyl orange, solutions (RE 17) | <i>E</i> | 266 |
| 2-Methylphenoxyacetic acid (RE 67) | <i>E</i> | 334 |
| Methyl red, solutions (RE 18) | <i>E</i> | 266 |
| Methyl undec-10-enoate (RE 89.2) | <i>E</i> | 361 |
| Metolachlor [400], method | <i>D</i> | 134 |
| Metribuzin [283], method | <i>D</i> | 137 |
| Metsulfuron [441], method, TC, WG | <i>H</i> | 204 |
| WP | <i>K</i> | 95 |
| Mevinphos [45] | <i>H</i> | 212 |
| MGK 264 + pyrethrum + piperonylbutoxide [345 + 32 + 33], method | <i>H</i> | 239 |
| Mineral acidity of dinitro compounds (MT 109) | <i>F</i> | 263 |
| Miscibility with hydrocarbon oil (MT 23) | <i>F</i> | 85 |
| Mineral oil (RE 36) | <i>E</i> | 292 |

CUMULATIVE INDEX

| | CIPAC | page |
|---|-----------------|------------------|
| Molinate [235], method | <i>IB</i> | <i>1866</i> |
| Monocrotophos [287], method | <i>E</i> | <i>145</i> |
| Mordant black, solutions (RE 53) | <i>E</i> | <i>316</i> |
| | | |
| Nabam [21], method | <i>E</i> | <i>151</i> |
| Naphtholbenzein, solution (RE 54) | <i>E</i> | <i>317</i> |
| Neutrality (MT 74) | <i>F</i> | <i>204</i> |
| Neutral oil, determination of distillation range (MT 61) | <i>F</i> | <i>183</i> |
| Niclosamide [599], method, TC, WP, EC | <i>J</i> | <i>84</i> |
| SC | <i>K</i> | <i>96</i> |
| Nicosulfuron [709], method | <i>M</i> | <i>121</i> |
| OD | <i>O</i> | <i>92</i> |
| Nitric acid, solutions (RE 51) | <i>E</i> | <i>313</i> |
| Nitromethane (RE 42) | <i>E</i> | <i>303</i> |
| Nitron (RE 78) | <i>E</i> | <i>345</i> |
| Nitron complexes, preparation of (MT 105) | <i>F</i> | <i>259</i> |
| Nitrophenols, determination of (MT 133) | <i>F</i> | <i>327</i> |
| Nitrothal-isopropyl [382], method | <i>E</i> | <i>154</i> |
| Novaluron [672], method | <i>L</i> | <i>91</i> |
| | | |
| Octacosane (RE 145) | <i>E</i> | <i>381</i> |
| Oil content of emulsifiable concentrates (MT 146) | <i>F</i> | <i>345</i> |
| Oil insoluble material (MT 35) | <i>F</i> | <i>108</i> |
| Omethoate [202], method | <i>E</i> | <i>159</i> |
| Organic chlorine, | | |
| potassium-xylene method (MT 38.1) | <i>F</i> | <i>118</i> |
| Stepanov method (MT 38.2) | <i>F</i> | <i>120</i> |
| semi-micro determination of total chlorine (MT 38.3) | <i>F</i> | <i>121</i> |
| Organochlorine pesticides, determination in aqueous emulsions (MT 65) | <i>F</i> | <i>190</i> |
| Oxadixyl [397], method | <i>G</i> | <i>122</i> |
| Oxamyl [342], method | <i>M</i> | <i>129</i> |
| Oxydemeton-methyl [171], method, HPLC | <i>E</i> | <i>163</i> |
| TLC | <i>IB</i> | <i>1871</i> |
| HPLC | <i>K</i> | <i>97</i> |
| | | |
| Packing of columns for gaschromatography (MT 149) | <i>F</i> | <i>349</i> |
| Palmitic acid (RE 80) | <i>E</i> | <i>348</i> |
| Paraffin, liquid (RE 8) | <i>E</i> | <i>255</i> |
| Paraquat [56], aqueous solutions, method | <i>E</i> | <i>166</i> |
| identity tests | <i>G</i> | <i>128</i> |
| free 4,4'-bipyridyl | <i>H</i> | <i>216</i> |

CUMULATIVE INDEX

| | CIPAC | page |
|---|-----------------|------------|
| terpyridines in | <i>M</i> | 141 |
| Paraquat + diquat [55 + 56], method | <i>E</i> | 75 |
| Paraquat dichloride (PP 55) | <i>E</i> | 390 |
| Parathion [10], method, GC | <i>1B</i> | 1875 |
| HPLC | <i>1B</i> | 1876 |
| Parathion-methyl [487], method, GC, TC, WP, EC | <i>1B</i> | 1879 |
| HPLCTC,EC | <i>1B</i> | 1881 |
| HPLCCS | <i>1C</i> | 2169 |
| GC OL,DP | <i>E</i> | 171 |
| HPLCOL | <i>E</i> | 171 |
| HPLC, determination of S-methyl isomer in TC | <i>E</i> | 169 |
| HPLC, determination of S-methyl isomer in SC | <i>E</i> | 172 |
| Parathion-methyl, free, determination in CS formulations (MT 188) | <i>L</i> | 132 |
| Particle size analysis by laser diffraction (MT 187) | <i>K</i> | 153 |
| Particle size, formulations without carriers (MT 42.1) | <i>F</i> | 133 |
| formulations containing carriers (MT 42.2) | <i>F</i> | 138 |
| Particle size distribution of a DDT dispersible powder (MT 43) | <i>F</i> | 140 |
| PCAF Reagent (RE 146) | <i>E</i> | 382 |
| Pebulate [236], method | <i>1B</i> | 1887 |
| Pencycuron [402], method | <i>E</i> | 173 |
| Pendimethalin [357], method | <i>M</i> | 148 |
| Pentacyanoaminoferroate (RE 146) | <i>E</i> | 382 |
| Perchloric acid, standard solutions (RE 43) | <i>E</i> | 304 |
| Permethrin, identity test (MT 163) | <i>F</i> | 404 |
| Permethrin [331], method, TC, WP, EC, WG, DP | <i>1C</i> | 2173 |
| TC, EW, LN | <i>M</i> | 155 |
| HPLC, enantiomeric purity in TC, EW | <i>O</i> | 94 |
| EC, LN, capillary GC + piperonyl butoxide | <i>O</i> | 97 |
| Persistent foaming (MT 47) | <i>F</i> | 152 |
| Persistent foam, general method (MT 47.3) | <i>O</i> | 177 |
| Petroleum spirit (RE 45) | <i>E</i> | 307 |
| pH, determination of (MT 75) | <i>F</i> | 205 |
| revised method (MT 75.3) | <i>J</i> | 131 |
| Phenmedipham [77], method, titration method | <i>1C</i> | 2181 |
| HPLC | <i>1C</i> | 2184 |
| HPLC (revised) | <i>J</i> | 92 |
| Phenolphthalein, solution (RE 19) | <i>E</i> | 267 |
| <i>d</i> -Phenothrin [777], method | <i>L</i> | 96 |
| Phenylurea herbicides, detection and identification of impurities in substituted (MT 142) | <i>F</i> | 336 |
| Phorate [173], method | <i>1B</i> | 1890 |
| Phosalone [109], method | <i>D</i> | 141 |

CUMULATIVE INDEX

| | CIPAC | page |
|---|--------------------|------------|
| Phosphamidon [110], method, method | <i>E</i> | 179 |
| Phosphate buffer solutions (MT 19) | <i>F</i> | 69 |
| Phosphates, test for (MT 120) | <i>F</i> | 285 |
| Phosphorus in active ingredients, spectrophotometric determination after TLC separation (MT 145) | <i>F</i> | 340 |
| Phosphorus pentoxide desiccant (MT 24) | <i>F</i> | 86 |
| Phoxim [364], method | <i>1C</i> | 2187 |
| Picloram [174], method (salts) | <i>1B</i> | 1893 |
| Picloram + 2,4-D [174 + 1] | <i>1B</i> | 1757 |
| HPLC, TC, SL | <i>L</i> | 104 |
| Piperonyl butoxide [33], method, capillary GC, TC | <i>O</i> | 104 |
| TC | <i>1C</i> | 2190 |
| EW | <i>M</i> | 163 |
| LN | <i>N</i> | 111 |
| LN + deltamethrin | <i>O</i> | 111 |
| LN + permethrin | <i>O</i> | 111 |
| + pyrethrum [33 + 32] | <i>1C</i> | 2209 |
| + pyrethrum + MGK 264 [33 + 32 + 345] | <i>H</i> | 239 |
| dihydrosafrole in, TC | <i>O</i> | 104 |
| Pirimicarb, identity test (MT 164) | <i>F</i> | 406 |
| Pirimicarb [231], method | <i>H</i> | 217 |
| Pirimiphos-ethyl, identity test (MT 164) | <i>F</i> | 406 |
| Pirimiphos-ethyl [492], method | <i>D</i> | 146 |
| Pirimiphos-methyl, identity test (MT 164) | <i>F</i> | 406 |
| Pirimiphos-methyl [239], method, GC, TC, WP, EC, UL, HN, DP | <i>1C</i> | 2192 |
| capillary method, GC, EC, CS | <i>O</i> | 112 |
| isopirimiphos-methyl in, TC, CS | <i>O</i> | 116 |
| <i>O,O</i>-dimethyl phosphorochloridothioate in, TC | <i>O</i> | 118 |
| <i>O,O,O</i>-trimethyl phosphorothioate in, TC | <i>O</i> | 118 |
| <i>O,O,S</i>-trimethyl phosphorothioate in, TC | <i>O</i> | 118 |
| <i>O,O,S</i>-trimethyl phosphorodithioate in, TC | <i>O</i> | 118 |
| Pirimiphos-methyl, free, determination in formulations (MT 189.2) | CS <i>O</i> | 193 |
| Pirimiphos-methyl, release rate of CS formulations (MT 190.2) | <i>O</i> | 199 |
| Pirimiphos-methyl, pure (PP 239.a) | <i>1C</i> | 2337 |
| Potassium bromate, standard solution (RE 41) | <i>E</i> | 303 |
| Potassium bromate/bromide, standard solutions (RE 82) | <i>E</i> | 351 |
| Potassium cyanate [197], method | <i>1C</i> | 2200 |
| Potassium dichromate, standard solutions (RE 20) | <i>E</i> | 267 |
| Potassium dihydrogen orthophosphate (RE 134) | <i>E</i> | 376 |
| Potassium hydroxide, standard solutions (RE 21) | <i>E</i> | 268 |

CUMULATIVE INDEX

| | CIPAC | page |
|--|-----------------|------------|
| Potassium iodate (RE 98) | <i>E</i> | 367 |
| Potassium iodate/iodide, standard solution (RE 15) | <i>E</i> | 264 |
| Potassium permanganate, standard solution (RE 22) | <i>E</i> | 270 |
| Potassium plumbite (RE 75) | <i>E</i> | 343 |
| Potassium thiocyanate, standard solution (RE 23) | <i>E</i> | 271 |
| Pour and tap bulk density of granular materials (MT 159) | <i>F</i> | 390 |
| Pourability of suspension concentrates (MT 148) | <i>F</i> | 348 |
| revised method (MT 148.1) | <i>J</i> | 133 |
| Pour point of mineral oil (MT 139) | <i>F</i> | 331 |
| Prallethrin [743], method | <i>L</i> | 110 |
| Prochloraz [407], method, TC | <i>M</i> | 165 |
| EC | <i>N</i> | 121 |
| Prochloraz-ZnCl ₂ Complex TC | <i>N</i> | 119 |
| Procymidon [383], method | <i>K</i> | 105 |
| Profenophos; [461], method | <i>H</i> | 222 |
| Prometon [175], method, titration method | <i>1C</i> | 2202 |
| GLC | <i>1C</i> | 2203 |
| Prometryn [93], method | <i>H</i> | 228 |
| Propachlor [176], method | <i>D</i> | 151 |
| Propamocarb [399] hydrochloride, method | <i>E</i> | 184 |
| Propan-2-ol (RE 40) | <i>E</i> | 302 |
| Propazine [92], method | <i>H</i> | 232 |
| Propetamphos [394], method | <i>E</i> | 187 |
| Propham + chlorpropham [63 + 43] | <i>1C</i> | 2025 |
| Propiconazole [408], method | <i>G</i> | 129 |
| Propineb [177], method | <i>H</i> | 236 |
| Propoxur [80], method | <i>D</i> | 155 |
| Provisional CIPAC method | <i>O</i> | 1 |
| Pure pesticides | <i>D</i> | 186 |
| Pyraclostrobin [657], method | <i>M</i> | 170 |
| Pyraoxystrobin [964], method, TC, SC | <i>O</i> | 134 |
| Pyrazophos [350], method | <i>1C</i> | 2206 |
| Pyrethrum [32], method | <i>1C</i> | 2209 |
| Pyrethrum + piperonylbutoxide [32 + 33] | <i>1C</i> | 2209 |
| Pyrethrum + MGK 264 + piperonylbutoxide [32 + 345 + 33] | <i>H</i> | 239 |
| Pyridaben [583], method | <i>J</i> | 94 |
| Pyridate [447], method | <i>G</i> | 137 |
| Pyridine (RE 48) | <i>E</i> | 310 |
| Pyriproxyfen [715], method | <i>M</i> | 178 |
| LN formulation | <i>O</i> | 143 |

CUMULATIVE INDEX

| | CIPAC | page |
|--|-----------------|------------|
| Qualitative procedure for confirmation of the presence of dithio-carbamates or thiuram disulfide (MT 153) | <i>F</i> | 360 |
| Quaternary ammonium compounds [MT 199], method | <i>O</i> | 223 |
| Quinclorac [493], method | <i>H</i> | 244 |
| Quinmerac [563], method | <i>J</i> | 101 |
| Quintozene [78], method | <i>IC</i> | 2213 |
| Rate of solution of the sodium salts of phenoxyalkanoic acid herbicides and their solid formulations (MT 60) | <i>F</i> | 182 |
| Referee method | <i>O</i> | 1 |
| Removal of dyes (MT 45) | <i>F</i> | 147 |
| Repeatability | <i>O</i> | 3 |
| Reproducibility | <i>O</i> | 3 |
| Residue on evaporation, low boiling products (MT 80.1) | <i>F</i> | 213 |
| cresols (MT 80.2) | <i>F</i> | 214 |
| Rimsulfuron [716], method | <i>M</i> | 189 |
| Retention test of seed treatment powders (MT 147) | <i>F</i> | 346 |
| Rotenone [38], method | <i>IC</i> | 2217 |
| Sampling of water dispersible granules (MT 166) | <i>F</i> | 415 |
| Sand for germination tests (MT 25) | <i>F</i> | 86 |
| Seed adhesion test for powders for seed treatment (MT 83) | <i>F</i> | 216 |
| Seed treatment formulations, liquid, determination of seed-seed uniformity of distribution (MT 175) | <i>F</i> | 438 |
| Separation and identification of herbicides (MT 97) | <i>F</i> | 241 |
| Sethoxydim [401], method | <i>E</i> | 193 |
| Sieve analysis, | | |
| dry sieving-dusts (MT 59.1) | <i>F</i> | 177 |
| dry sieving-granular products (MT 59.2) | <i>F</i> | 179 |
| dry sieving-water dispersible granules (MT 170) | <i>F</i> | 420 |
| sieve test for granular material (MT 59.4) | <i>F</i> | 179 |
| wet sieving (MT 59.3) | <i>F</i> | 135 |
| wet sieving using recycled water (MT 182) | <i>J</i> | 416 |
| wet sieving-water dispersible granules (MT 167) | <i>F</i> | |
| wet sieve test (MT 185) | <i>K</i> | 149 |
| Silanisation of gas chromatographic columns (MT 113) | <i>F</i> | 277 |
| Silica for chromatography, | | |
| silica (MT 21.1) | <i>F</i> | 71 |
| Sorbisil M 60 (MT 21.2) | <i>F</i> | 72 |
| Florisil (MT 21.3) | <i>F</i> | 73 |
| Silicates, test for (MT 121) | <i>F</i> | 287 |
| Silver diethyldithiocarbamate (RE 44) | <i>E</i> | 306 |

CUMULATIVE INDEX

| | CIPAC | page |
|---|-----------------|------------|
| Silver nitrate, standard solutions (RE 24) | <i>E</i> | 273 |
| Simazine [22], method, | <i>H</i> | 250 |
| Simetryn [179], method, titration method | <i>1B</i> | 1895 |
| GLC | <i>1B</i> | 1896 |
| Sodium anilinobenzene sulfonate (RE 147) | <i>E</i> | 383 |
| Sodium biphenyl (RE 46) | <i>E</i> | 308 |
| Sodium bis(2-ethylhexylcarbonyl) ethane sulfonate (RE 71) | <i>E</i> | 338 |
| Sodium bis(2-ethylhexyl) sulfosuccinate (RE 71) | <i>E</i> | 338 |
| Sodium carbonate (RE 52) | <i>E</i> | 315 |
| Sodium chlorate [7], method | <i>E</i> | 197 |
| Sodium diethyldithiocarbamate trihydrate (RE 96) | <i>E</i> | 365 |
| diSodium dihydrogen ethylenediaminetetra-acetate (RE 37) | <i>E</i> | 295 |
| Sodium diphenylaminesulfonate (RE 147) | <i>E</i> | 383 |
| Sodium hydroxide, standard solutions (RE 25) | <i>E</i> | 273 |
| Sodium nitrite, standard solutions (RE 39) | <i>E</i> | 299 |
| Sodium pentacyanonitrosylferrate(III) (RE 146) | <i>E</i> | 382 |
| diSodium tetraborate (RE 133) | <i>E</i> | 376 |
| Sodium thiosulfate, standard solutions (RE 26) | <i>E</i> | 276 |
| Solubility in acetone (MT 5) | <i>F</i> | 21 |
| Solubility in ethanol (MT 7), | <i>F</i> | 23 |
| Solubility in hexane (MT 6), | <i>F</i> | 22 |
| Solubility in organic solvents (MT 181) | <i>H</i> | 314 |
| Solubility in sodium hydroxide, | | |
| phenoxyalkanoic herbicides (MT 71.1) | <i>F</i> | 200 |
| cresols (MT 71.2) | <i>F</i> | 200 |
| bromoxynil and ioxynil (MT 71.3) | <i>F</i> | 201 |
| Solubility in triethanolamine (MT 76) | <i>F</i> | 207 |
| Solubility of salts and alkali insoluble-dinitro compounds (MT 108) | <i>F</i> | 262 |
| Soluble alkalinity (MT 81) | <i>F</i> | 215 |
| Soluble chlorides (MT 82) | <i>F</i> | 215 |
| Solution properties of water soluble tablets (MT 196) | <i>O</i> | 210 |
| Solvent naphtha, aromatic (RE 62) | <i>E</i> | 328 |
| Specific gravity and density, | | |
| hydrometer method (MT 3.1) | <i>F</i> | 11 |
| pycnometer method (MT 3.2) | <i>F</i> | 13 |
| density of suspension concentrates (MT 3.3) | <i>F</i> | 18 |
| Spinosad [636], method, TC, SC, GR | <i>L</i> | 121 |
| DT | <i>M</i> | 197 |
| EC, GR | <i>O</i> | 146 |
| Spontaneity of dispersion, of suspension concentrates (MT 160) | <i>F</i> | 391 |
| of water dispersible granules (MT 174) | <i>F</i> | 435 |

CUMULATIVE INDEX

| | CIPAC | page |
|---|-----------|------------|
| Stability of diluted oil products, | <i>F</i> | 167 |
| tar-petroleum oil (MT 55.1) | <i>F</i> | 168 |
| petroleum and DNOC petroleum oils (MT 55.2) | <i>F</i> | 168 |
| petroleum oils for orchard use (MT 55.3) | <i>F</i> | 168 |
| petroleum oils for glasshouse use (MT 55.4) | | |
| Stability of diluted tar and petroleum oil products, | | |
| tar-oils-miscible and stock emulsion types (MT 49.1) | <i>F</i> | 155 |
| petroleum oil-miscible type (MT 49.2) | <i>F</i> | 156 |
| Stability of diluted and petroleum oil concentrate products, | | |
| miscible type (MT 52.1) | <i>F</i> | 159 |
| Stability of undiluted DNOC-petroleum, petroleum and tar- | | |
| petroleum oil products (MT 54) | <i>F</i> | 167 |
| Stability of undiluted tar oil products, | | |
| miscible type (MT 48.1) | <i>F</i> | 154 |
| stock emulsion type (MT 48.2) | <i>F</i> | 154 |
| Stability of tar-petroleum and petroleum oil products-undiluted, | | |
| miscible type (MT 51.1) | <i>F</i> | 158 |
| Standard waters, | | |
| CIPAC standard waters A - G (MT 18.1) | <i>F</i> | 59 |
| H and J (MT 18.2) | <i>F</i> | 63 |
| non-CIPAC standard waters (MT 18.3) | <i>F</i> | 64 |
| simplified procedure (MT 18.5) | <i>H</i> | 302 |
| Starch solution (RE 27) | <i>E</i> | 280 |
| Starch, indicator paper (RE 32) | <i>E</i> | 284 |
| Sulfated ash, general method (MT 29) | <i>F</i> | 91 |
| Sulfometuron-methyl [610], method | <i>J</i> | 108 |
| Sulfotep [198], method | <i>H</i> | 255 |
| Sulfur [18], method | <i>E</i> | 203 |
| Sulfuric acid, standard solutions (RE 28) | <i>E</i> | 280 |
| Suspensibility of formulations forming suspensions on dilution with water (MT 184) | <i>K</i> | 142 |
| Suspensibility of aqueous suspension concentrates (MT 161) | <i>F</i> | 394 |
| Suspensibility of water dispersible granules (MT 168) | <i>F</i> | 417 |
| Suspensibility of water dispersible powders, | | |
| CIPAC method (MT 15.1) | <i>F</i> | 45 |
| AID method (MT 15.2) | <i>F</i> | 48 |
| simplified CIPAC method (MT 177) | <i>F</i> | 445 |
| Suspo-emulsions, dispersion stability (MT 180) | <i>H</i> | 310 |
| SE, CD and OD formulations, (MT 180) | O | 192 |
| 2,4,5-T [6], method, titration method | <i>1C</i> | 2220 |
| GC | <i>1C</i> | 2221 |

CUMULATIVE INDEX

| | CIPAC | page |
|---|-----------------|------------|
| HPLC | <i>IC</i> | 2221 |
| - salts | <i>IC</i> | 2223 |
| -esters | <i>IC</i> | 2223 |
| 2,4,5-T + 2,4-D [6 + 1] | <i>IC</i> | 2072 |
| 2,4,5-T, pure (PP 6) | <i>IC</i> | 2312 |
| Tannic acid (RE 143) | <i>E</i> | 376 |
| Tap density, of powders (MT 33) | <i>F</i> | 104 |
| of water dispersible granules (MT 169) | <i>F</i> | 418 |
| 2,3,6-TBA [49], method titration method | <i>IC</i> | 2225 |
| GLC | <i>IC</i> | 2227 |
| Na-salt titration method | <i>IC</i> | 2228 |
| GLC | <i>IC</i> | 2229 |
| 2,3,6-TBA + MCPA(salt) [49 + 2] | <i>IC</i> | 2148 |
| 2,3,6-TBA, pure (PP49) | <i>IC</i> | 2314 |
| TCDD, determination of | | |
| in 2,4,5-T technical (MT 151.1) | <i>F</i> | 352 |
| in 2,4,5-T technical esters (MT 151.2) | <i>F</i> | 355 |
| Tebuconazole [494], method | <i>H</i> | 261 |
| Temephos [340], method | <i>IC</i> | 2230 |
| Tentative CIPAC method | <i>O</i> | 1 |
| Terbufos [459], method | <i>H</i> | 269 |
| Terbutylazine [234], method, titration method | <i>IB</i> | 1897 |
| GLC | <i>IB</i> | 1898 |
| Terbutryn [212], method | <i>H</i> | 275 |
| Tetradifon [113], method | <i>IB</i> | 1901 |
| Thallium [181], method | <i>IC</i> | 2234 |
| Thiabendazole [323], method | <i>E</i> | 211 |
| Thiacloprid [631], method | <i>M</i> | 198 |
| Thifensulfuron [452], method | <i>K</i> | 114 |
| Thiobencarb [388], method | <i>D</i> | 159 |
| Thiodicarb [543], method | <i>H</i> | 279 |
| Thiomethoxam [636], method, TC, WG, SC, FS | <i>O</i> | 147 |
| Thiometon [115], infrared method | <i>H</i> | 285 |
| gas chromatographic method | <i>E</i> | 218 |
| Thiophanate-methyl [362], method | <i>D</i> | 162 |
| Thiram [24], method | <i>D</i> | 169 |
| Thiram, qualitative procedure for confirmation of the presence of (MT 153) | <i>F</i> | 360 |
| Thymol blue, solutions (RE 30) | <i>E</i> | 283 |
| Thymolphthalein, solution (RE 30) | <i>E</i> | 344 |
| Titanium(III)chloride (RE 83) | <i>E</i> | 352 |

CUMULATIVE INDEX

| | CIPAC | page |
|--|-----------------|------------|
| Titanium(III)sulfate (RE 85) | <i>E</i> | 358 |
| TMPS, method | <i>E</i> | 142 |
| <i>i</i> -TMPS, method | <i>E</i> | 142 |
| Toluene (RE 152) | <i>E</i> | 384 |
| Toluene solubles (MT 90) | <i>F</i> | 222 |
| Toluene, method (MT 198) | <i>O</i> | 215 |
| Tolyfluanid [275], method | <i>E</i> | 222 |
| <i>o</i> -Tolyloxyacetic acid (RE 67) | <i>E</i> | 334 |
| Total chlorides (MT 100) | <i>F</i> | 254 |
| Total hardness of water (MT 73.1) | <i>O</i> | 179 |
| Transfluthrin [741], method, TC, SL | <i>K</i> | 121 |
| TC, VL | <i>L</i> | 128 |
| identification of isomers, TC | <i>O</i> | 158 |
| Triadimefon [352], method | <i>1C</i> | 2236 |
| revised method, TC, WP, EC, GR | <i>N</i> | 126 |
| Triadimenol [398], method | <i>E</i> | 224 |
| revised method, TC, WP, WG EC, SC, SE, EW | <i>N</i> | 134 |
| Triazophos [353], method | <i>H</i> | 288 |
| Tribenuron -methyl [546], method | <i>K</i> | 128 |
| 1,1,1-Trichloroethane (RE 95) | <i>E</i> | 365 |
| Trifloxystrobin [617], method, TC, AL, EC, FS, SC, WG | <i>O</i> | 161 |
| Triflumuron [548], method | <i>J</i> | 114 |
| Trifluralin [183], method | <i>H</i> | 292 |
| Triforine [360], method | <i>H</i> | 299 |
| <i>O,O,S</i> -Trimethylphosphorodithioate (RE 87) | <i>E</i> | 360 |
| in pirimiphos-methyl, TC | <i>O</i> | 118 |
| <i>O,O,O</i> -Trimethylphosphorothioate, method | <i>E</i> | 142 |
| in pirimiphos-methyl, TC | <i>O</i> | 118 |
| <i>O,O,S</i> -Trimethylphosphorothioate, method | <i>E</i> | 142 |
| in pirimiphos-methyl, TC | <i>O</i> | 118 |
| (Tris(1,10-phenanthroline)- <i>N,N'</i>)iron(II) sulfate (RE 149) | <i>E</i> | 383 |
| Ultraviolet absorption test for ethylenebisdithiocarbamates (MT 165) | <i>F</i> | 411 |
| Undec-10-enoic acid (RE 89) | <i>E</i> | 360 |
| Unulfonated residue of neutral oil (MT 57) | <i>F</i> | 170 |
| Urea herbicides, identification of (MT 137) | <i>F</i> | 329 |
| Vernolate [237], method | <i>1B</i> | 1905 |
| Viscosity, viscosity of transparent and opaque liquids (MT 22.1) | <i>F</i> | 75 |
| viscosity Redwood (MT 22.2) | <i>F</i> | 81 |
| viscosity of mineral oil (MT 22.3) | <i>F</i> | 83 |

CUMULATIVE INDEX

| | CIPAC | page |
|---|-----------------|------------|
| Viscosity of liquids by rotational viscometry (MT 192) | <i>L</i> | 145 |
| revised method | <i>M</i> | 210 |
| Vinclozolin [280], method | <i>D</i> | 173 |
| Volatility of neutral oil, | | |
| preliminary examination (MT 56.1) | <i>F</i> | 170 |
| full method (MT 56.2) | <i>F</i> | 170 |
| Wash resistance index of LN's (MT 195) | <i>O</i> | 205 |
| Water, for laboratory use (RE 130) | <i>E</i> | 374 |
| Water, | | |
| Karl Fischer method | <i>F</i> | 92 |
| Dean and Stark method (MT 30.2) | <i>F</i> | 94 |
| free water 'speedy' method (MT 30.3) | <i>F</i> | 95 |
| water in acetone solutions (MT 30.4) | <i>F</i> | 96 |
| Karl Fischer method using pyridine-free reagents (MT 30.5) | <i>J</i> | 120 |
| Water content and suspended solids, | | |
| water content (MT 40.1) | <i>F</i> | 131 |
| suspended solids (MT 40.2) | <i>F</i> | 131 |
| Water insoluble material, | | |
| hot solution of the sample (MT 10.1) | <i>F</i> | 27 |
| cold solution of the sample (MT 10.2) | <i>F</i> | 27 |
| coarse material-water insoluble (MT 10.3) | <i>F</i> | 28 |
| insoluble material in pesticide aqueous solutions (MT 10.4) | <i>F</i> | 29 |
| Water solubility, | | |
| column elution method (MT 157.2) | <i>F</i> | 380 |
| flask method (MT 157.3) | <i>F</i> | 384 |
| preliminary test (MT 157.1) | <i>F</i> | 379 |
| Water soluble bags, dissolution rate of (MT 176) | <i>F</i> | 440 |
| Water soluble granules, degree of dissolution and solution stability (MT 179) | <i>H</i> | 307 |
| Dissolution degree and solution stability (MT 179.1) | <i>O</i> | 189 |
| replaces MT 179 | | |
| Water soluble material (MT 9) | <i>F</i> | 26 |
| Wettability (MT 53) | <i>F</i> | 160 |
| Xylene (RE 10) | <i>E</i> | 256 |
| Xylene insoluble material (MT 11) | <i>F</i> | 30 |
| Xylenol Orange (RE 93) | <i>E</i> | 364 |
| Zinc amalgam (RE 33) | <i>E</i> | 284 |
| Zinc chloride, solutions (RE 94) | <i>E</i> | 364 |

CUMULATIVE INDEX

| | CIPAC | page |
|-------------------------------------|----------|------|
| Zinc, in dithiocarbamates (MT 94.1) | <i>F</i> | 235 |
| Zinc, in zineb, method | <i>E</i> | 231 |
| Zinc, in ziram, method | <i>E</i> | 235 |
| Zineb [25], method | <i>E</i> | 230 |
| Zineb + copper [25 + 44], method | <i>H</i> | 100 |
| Ziram [31], method | <i>E</i> | 236 |
| Ziram + copper [31 + 44], method | <i>H</i> | 101 |