


# CIPAC STATUS REPORT

01/07/2005



## 0609 Epoxiconazole

---

Allocated to D

CIPAC methods published in :

CIPAC K, p. 38

**CIPAC** 42th meeting, July 1998 in York

A small scale study with a HPLC method has been started within DAPA.

**CIPAC** 42nd meeting, July 1998 in York

A small scale study with a HPLC method has been started within DAPA.

**CIPAC** 43rd meeting, June 1999 in Budapest

Mr. Pawliczek presented the report of a small scale study performed by DAPA. It was a study with a capillary GLC method on 1 TC and 2 SC's, one of them with a second a.i. The replacement of He by N<sub>2</sub> had not been tested. It was remarked that an identity test might be necessary to determine whether or not it is a mixture of the four diastereomers. The results of the study were good, a CIPAC study will be started soon.

**CIPAC** 44th meeting, June 2000 in Granada

Mr. Pawliczek presented the report of a CIPAC study (CIPAC/4190). It was a study with a capillary GLC method on 1 TC and 2 SC's, 1 EC and 1 SE. The results obtained from the study were good. Decision The capillary GC method for epoxiconazole technical and formulations (EC, SC, SE), CIPAC/4189, has been accepted as provisional CIPAC method.

**CIPAC** 45th meeting, June 2001 in Bangkok

Decision The provisional capillary GC method for epoxiconazole technical and formulations (EC, SC, SE), CIPAC/4189, has been accepted as full CIPAC method