

Order form for CIPAC pre-published methods

Adopted CIPAC methods are available on a per-method basis, as long as they are not yet published in a Handbook. CIPAC charges a nominal fee per method.

Instruction

The list of the pre-published methods below is exhaustive.

Tick the method or methods you wish to receive. The methods are sent as pdf files attached to an e-mail – a working e-mail address is therefore mandatory.

Each method is charged with GBP 5. Please sum the number of methods you wish to receive and calculate the grand total. Send the form duly filled and signed to Marston Book Services (see details below) by fax.

As soon as your order is processed, you shall receive the pdf files by e-mail attachment.

Terms of use

By ordering one or more of these methods, the user accepts the conditions of use as follows:

- these are draft methods and are subject to revision without further notice
- these methods will appear in revised and edited form in CIPAC Handbook O presumably in the year 2016.
- The CIPAC copyright and disclaimer (cf. CIPAC Handbook N, page ii) apply to all pre-published methods

CIPAC No	Name	Description	Order (tick)
789	Amisulbrom	Reversed phase HPLC method (CIPAC/4883) for the determination of amisulbrom in TC, WG and SC formulations (ISBN 90295184X).	<input type="radio"/>
570	Chlorfenapyr	Reversed phase HPLC method (CIPAC/4826) for the determination of chlorfenapyr in TC and SC (ISBN 902951912).	<input type="radio"/>
653	Cyazofamid	Reversed phase HPLC method (CIPAC/4833) for the determination of cyazofamid in TC and SC (ISBN 902951939).	<input type="radio"/>
333	Deltamethrin	Enantioselective HPLC method (CIPAC/4907) for the determination of the enantiomeric ratio of deltamethrin and its stereoisomers not belonging to the active ingredient definition in TC (ISBN 902951874).	<input type="radio"/>
739	Dimoxystrobin	Capillary GC method (CIPAC/4710) for determination of dimoxystrobin content in TC, SE and SC formulations (ISBN 902951602).	<input type="radio"/>
595	Flazasulfuron	Reversed phase HPLC method (CIPAC/4831) for the determination of flazasulfuron in TC and WG (ISBN 902951920).	<input type="radio"/>
521	Fluazinam	Reversed phase HPLC method (CIPAC/4727) for the determination of fluazinam in TC and SC formulation (ISBN 902951696)	<input type="radio"/>
578	Flumioxazin	Reversed phase HPLC method (CIPAC/4763) for the determination of flumioxazin in TC and WP formulations (ISBN 902951629)	<input type="radio"/>
585	Fosthiazate	Reversed phase HPLC method (CIPAC/4829) for the determination of fosthiazate in TC and GR (ISBN 902951793).	<input type="radio"/>
709	Nicosulfuron	Extension of the scope (CIPAC/4903) of CIPAC method 709/TC/M/3 for the determination of the nicosulfuron content of oil-based suspension concentrate formulations (OD) (ISBN 902951882)	<input type="radio"/>

331	Permethrin	Method extension to the capillary GC method 331/TC/M/3 (CIPAC/4719) for the determination of the content of permethrin in EC formulations (ISBN 902951319).	○
33	Piperonyl butoxide	Capillary GC method (CIPAC/4765) for the determination of the content of piperonyl butoxide in TC (ISBN 902951815).	○
239	Pirimiphos-methyl	Capillary GC-FID method (CIPAC/4963) for the determination of total pirimiphos-methyl in TC, EC and CS formulations (ISBN 902951645). Revision 2	○
715	Pyriproxyfen	Extension of the scope (CIPAC/4887) of CIPAC method 715/TC/M/2 for the determination of the pyriproxyfen content of a long lasting insecticidal mosquito net (incorporated type) (LN) containing permethrin and pyriproxyfen (ISBN 902951904).	○
636	Spinosad	Method extension to the reversed phase HPLC method 636/TC/M/3 (CIPAC/4721) for the determination of the content of spinosad in EC formulations (ISBN 902951335).	○
636	Spinosad	Method extension to the reversed phase HPLC method 636/TC/M/3 (CIPAC/4847) for the determination of the content of spinosad in GR formulations with lower content (ISBN 902951742).	○
637	Thiamethoxam	Reversed phase HPLC method (CIPAC/4845) for the determination of thiamethoxam in TC, WG, SC and FS (ISBN 902951785).	○
370	Brodifacoum	The reversed phase HPLC method (CIPAC/4942) for the determination of brodifacoum in TC and RB formulations was accepted as a provisional CIPAC method with the proposal of updating the description of the method concerning standard and sample preparation (ISBN 090295198X).	
964	Pyraoxystrobin	The reversed phase HPLC method (CIPAC/4936) for the determination of pyraoxystrobin in TC and SC formulations was accepted as a provisional CIPAC method with the proposal of amending the description of the method concerning sonication time and clarification of a possible inclusion of a note concerning addition of water for the sample preparation of the SC (ISBN 978-1-911009-51-1).	
617	Trifloxystrobin	The reversed phase HPLC method (CIPAC/4954) for the determination of trifloxystrobin in TC, EC, FS, SC, WG and AL formulations was accepted as a provisional CIPAC method with the proposal of amending the description of the method concerning sonication time and clarification of a possible inclusion of a note concerning injection volume (ISBN 978-1-911009-02-3).	
331	Permethrin	The chiral HPLC method (CIPAC/4946) for the determination of the enantiomeric ratio of the four permethrin stereoisomers in technical active substance and EW formulation was accepted as an enantioselective identity test (ISBN 978-1-911009-03-0).	
741	Transfluthrin	The chiral HPLC method (CIPAC/4948) for the determination of the enantiomeric ratio of the four transfluthrin stereoisomers in technical active substance was accepted as an enantioselective identity test (ISBN 0902951998).	
635	Silthiofam	The reversed phase HPLC method using internal standard (CIPAC/5004) for the determination of silthiofam in TC and FS formulations was accepted as a provisional CIPAC method with the inclusion of a note that the addition of the internal standard can be done also volumetrically (ISBN 978-1-911009-05-4).	

715	Pyriproxyfen	The extension of the scope (CIPAC/4997) of CIPAC method 715/TC/M/3 for the determination of the pyriproxyfen content of a matrix release formulation (MR) (incorporated type) was accepted as provisional CIPAC method (ISBN 978-1-911009-06-1).	
794	Chlorantraniliprole	The reversed phase HPLC method using internal standard (CIPAC/5034) for the determination of chlorantraniliprole in TC, FS, WG and SC formulations was accepted as a provisional CIPAC method (ISBN 978-1-911009-09-2).	
693	Fenazaquin	The reversed phase HPLC method using external standardisation (CIPAC/5036) for the determination of fenazaquin in TC and SC formulations was accepted as a provisional CIPAC method (ISBN 978-1-911009-10-8).	
454	Alpha-cypermethrin	The extension of the scope (CIPAC/5043) of CIPAC method 454/LN/M/3.2 for the determination of the alpha-cypermethrin content of the long lasting insecticidal mosquito net (incorporated type) (LN) containing alpha-cypermethrin and pyriproxyfen, with the modification of having di-cyclohexyl phthalate as internal standard, instead of dioctyl phthalate, was accepted as a tentative CIPAC method, with the need for the provision of a second data set according to the provisions of the CIPAC guideline (ISBN 978-1-911009-11-5).	
738	Clothianidin	The extension of the scope (CIPAC/5051) of CIPAC method 738/WG/M/ for the determination of the clothianidin content of WP formulations was accepted as a provisional CIPAC method (ISBN 978-1-911009-13-9).	
MT 46.3	Accelerated storage Procedure	The extension of the method MT 46.3 for the accelerated storage procedure of the LN formulations regarding determination of active ingredient content and retention index was accepted as a provisional CIPAC MT method (ISBN 978-1-911009-04-7).	
MT 46.3	Accelerated storage Procedure	The extension of the scope (CIPAC/5045) of CIPAC method MT 46.3 for the accelerated storage procedure of the MR formulations regarding determination of active ingredient content and retention index was accepted as a tentative CIPAC MT method (ISBN 978-1-911009-12-2).	
MT 47.3	Persistent foam	Submethod using a standard measuring cylinder (BS EN ISO 4788:2005) (ISBN 902951831).	○
MT 73.1	Total hardness of water	Method for determination of total hardness of water using commercial complexometry (CIPAC/4769) (ISBN 902951718).	○
MT 179.1	Degree of dissolution	The extension of the scope (CIPAC/4891) of CIPAC method MT 179 to water soluble formulations (such as SG, SP, SS) were accepted as a full CIPAC MT method and the changes proposed to method MT 179.1 were accepted. (ISBN 902951963).	○
MT 189.2	Pirimiphos-methyl	Method extension to the CIPAC method MT 189 (CIPAC/4785) for the determination of the content of non-encapsulated pirimiphos-methyl ("free" a.i.) in CS formulations (ISBN 902951653).	○
MT 190.2	Pirimiphos-methyl	Method extension to the CIPAC method MT 190 (CIPAC/4783) for the determination of the release rate of pirimiphos-methyl in CS formulations (ISBN 902951661). Revision 2	○
MT 195	Wash resistance index of LN	This method is intended for determination of the wash resistance index of long-lasting insecticidal net (LN). It is	○

		suitable for coated and incorporated types of LN, in combination with an adopted analytical method for the active ingredient and proper net type (ISBN 902951823).	
MT 196	Solution properties of ST formulations	Method for determination of the residue of soluble tablets retained on a sieve (CIPAC/4771) (ISBN 90295170X).	○
MT 197	Disintegration of tablets	Method for the determination of the disintegration of tablets which have to be dissolved or dispersed in water (ST, WT, ...) (CIPAC/4894) was accepted as a provisional CIPAC MT method (ISBN 902951971).	○
MT 198	Toluene	The headspace GC method (CIPAC/4944) for the determination of toluene in formulations was accepted as a provisional CIPAC MT method (ISBN 0902951947).	
MT 199	Quaternary ammonium compounds	The potentiometric titration method utilizing an ionic surfactant electrode (CIPAC/4965) for the determination of quaternary ammonium compounds in concentrated and ready-to-use (RTU) disinfectant formulations was accepted as a provisional CIPAC method (ISBN 978-1-911009-52-8).	
MT XXX	Retention properties of pyriproxyfen MR formulations	The method for the determination of retention properties of pyriproxyfen matrix release formulations (CIPAC/4999) was accepted as a tentative CIPAC MT method with the request of additional validation data (ISBN 978-1-911009-07-8).	
		Total number of pre-published methods	

To:
Marston Book Services Ltd,
160 Milton Park,
Abingdon,
Oxfordshire OX14 4SD
England.
Telephone for orders: +44 1235 465500
Telefax: +44 1235 465556
E-Mail: direct.orders@marston.co.uk

Total amount due

Number of pre-published methods x

£4.26 plus VAT £0.74 (UK/EEC customers) = gross £5.00: @ £.....**

Name:

Address:

.....

ZIP: City:

Country:

e-mail*:

Tel. No:

Fax No:

* e-mail address is mandatory. Orders cannot be processed without your working e-mail address.

** EEC customers – please enter your VAT number if registered.....

Payment instructions

Payment: either by credit card (Visa and MasterCard only) or by prepayment:

- Charge my Visa or MasterCard (fill the form below)
- Please send me a Pro Forma Invoice.
- I enclose a sterling cheque payable Marston Book Services Limited.
- I have made a payment, net of all bank charges, direct to
Barclays Bank plc, Oxford. Sort code 20 65 18.
Swift code BARC GB22
Iban no GB84 20651800636835
Account 0063685

Please delete as applicable

Payment by credit card (Visa and MasterCard only)

Customers are paying via credit/debit card need to either fax the order to us
+44 (0)1235 465592 or call us on **+44 (0)1235 465577**.

Card Company:

Name of cardholder:

Card No:

Three digit Card Security Code (on back of card).....

Expiry Date:

Signature of cardholder:

Signature

.....

**Prices correct 30 September 2015, but may be subject to change.
Prices include handling as e-mail attachment. **Visa and MasterCard** are accepted.*